

## Table of Contents

# PTSD: Specialized Care and Supportive Services



Stressful and traumatic events happen throughout life. Being frightened and having an added layer of tension while you process these emotions is a normal reaction. If you begin to have flashbacks, recurring memories or dreams, distressing thoughts, or physical symptoms of stress about the event, you may have [Post Traumatic Stress Disorder \(PTSD\)](#). Let's delve into understanding PTSD and explore various care options available to help you through this difficult time.

## Understanding PTSD

If you've experienced or witnessed a shocking or dangerous event where you thought your own or someone else's life was in danger, you're at risk for developing

PTSD.

## **What is PTSD?**

PTSD is a mental health disorder that can manifest in various symptoms, including flashbacks, nightmares, hypervigilance, and avoidance behaviors. PTSD can significantly impact an individual's daily life, relationships, and well-being. It's essential for those affected to seek support and treatment to manage symptoms effectively and improve their quality of life.

## **The Prevalence of PTSD**

As with many mental health challenges, it's hard to know exactly how many people are affected by PTSD. Many people don't understand the disorder and, therefore, don't seek support.

However, from what we do know and the numbers available from the National Center for PTSD, approximately [5% of the U.S. population](#) is affected annually, and [6% of the U.S. population](#) will be diagnosed with PTSD at some point in their lives.

PTSD can affect anyone regardless of age, gender, or race. However, some groups are at higher risk of developing PTSD. Due to the traumatic events that women are more likely to experience, such as sexual assault and domestic violence, [8% of women, compared to 4% of men](#), will be diagnosed with PTSD at some point in their lives. Also, [7% of Veterans](#) will be diagnosed with PTSD. This number increases depending on deployment and particular war zones. The number is highest in veterans of Operations Iraqi Freedom and Enduring Freedom with [29% experiencing PTSD](#), while [21% of Persian Gulf War veterans](#) experienced PTSD.

PTSD is a widespread mental health concern. Understanding the signs and symptoms is necessary to get the help you need, especially if you're in one of the risk factor groups discussed above.

## **Recognizing the Signs of PTSD**

Recognizing and understanding the symptoms of PTSD is a crucial step in supporting those on their path to healing. [Studies](#) show that early recognition is critical for better treatment outcomes and decreased long-term effects. Early intervention can make a difference, helping manage symptoms and improving overall well-being. Plus, it can stop things from getting worse or leading to other issues like substance

abuse.

The [National Institute of Mental Health \(NIMH\)](#) reports that for an adult to be diagnosed with PTSD, they must have experienced four symptoms once or twice throughout one month. These symptoms include:

- One re-experiencing symptom
- One avoidance symptom
- Two arousal and reactivity symptoms
- Two cognition and mood symptoms

### **Re-Experiencing Symptoms**

Imagine feeling like you're back amid a traumatic event, even when you're safe and sound in the present moment. Re-experiencing symptoms manifest as vivid nightmares or sudden flashbacks, where it feels like you're reliving the experience through your senses—seeing, hearing, even smelling what you did during the trauma. Triggers, from movie scenes to news reports or conversations about similar accidents, can lurk everywhere.

### **Avoidance Symptoms**

Picture going to great lengths to steer clear of anything that might trigger memories of the traumatic event—whether it's avoiding specific places, people, or activities or even suppressing thoughts and discussions about the event altogether. This avoidance can extend to social situations, leading to emotional numbness or detachment from others. Some individuals cope by keeping themselves excessively busy, filling every moment with activity to avoid confronting their emotions.

### **Arousal and Reactivity Symptoms**

Feelings of being stuck in a perpetual state of alertness and unease are commonplace for those experiencing PTSD. Imagine feeling constantly jittery, as if every sound or movement sets off a surge of adrenaline, making it hard to concentrate or find peace. Even the smallest surprises can provoke exaggerated responses, leaving you feeling constantly on edge and easily irritable.

### **Cognition and Mood Symptoms**

After enduring a traumatic event, you may experience negative thoughts about yourself and others while also feeling emotionally numb and disinterested in activities you once enjoyed. Memory gaps or difficulty discussing the trauma may compound feelings of confusion and detachment from reality. Trust in life and others may diminish, while feelings of guilt and shame related to the event can be overwhelming, clouding your sense of self-worth and contributing to a bleak outlook on the future.

## **Causes of PTSD**

PTSD can stem from a spectrum of traumatic experiences. Each individual's perception and response to trauma differs, influencing the development and manifestation of PTSD symptoms.

A [study](#) published in Neuroscience & Biobehavioral Reviews indicates that there are individual risk factors that may predispose some people, more than others, to develop PTSD. Some of those risk factors include:

- Intense or prolonged trauma
- Previous traumatic events
- Severity of trauma
- Occupational exposure
- Existing mental health conditions
- Substance misuse
- Limited support
- Family history of mental health concerns

Just as there are risk factors in the development of PTSD, the [NIMH](#) reports there are also resilience factors to decrease your chances of developing PTSD. Resilience factors are deeply rooted in acknowledging the event and seeking early support. Other factors include:

- Accepting support from friends and family
- Learning self-acceptance of your actions during the trauma
- Maintaining and having healthy coping strategies
- Educating yourself on triggers and practicing how to respond

Dispelling misconceptions surrounding PTSD is essential. While combat situations are commonly associated with the disorder, PTSD can arise from any trauma.

Recognizing this diversity is pivotal in providing comprehensive support and treatment to those affected by PTSD.

The following triggers may lead to the development of PTSD:

- Combat exposure
- Domestic violence
- Sexual trauma
- Physical assault
- Natural disasters
- Serious accidents
- Childhood abuse or neglect
- Medical trauma
- Witnessing trauma
- Acts of terrorism
- Sudden loss
- Prolonged stress
- Occupational exposure

### **Psychological Impact of PTSD**

PTSD deeply impacts how people feel, act, and behave by triggering intense emotions like fear, anxiety, and sadness. Intrusive thoughts and memories disrupt thinking patterns, leaving individuals feeling helpless and hopeless. Over time, those with PTSD may also [struggle with other mental health issues](#) like depression or anxiety, compounding the challenges they face.

Avoidance behaviors and emotional numbing brought on by PTSD can strain relationships, impair communication, and hinder intimacy. In the workplace, PTSD can diminish concentration, memory, and decision-making abilities, impacting productivity and job satisfaction. Overall, PTSD diminishes quality of life, with symptoms varying widely among individuals and often complicated by other conditions.

### **Physiological Effects of PTSD**

When you think of PTSD, you may only think of the mental health aspects, but when your mind is in a constant state of hyper-awareness and stress, your body is affected. There has been a considerable underestimation of the role of trauma –

PTSD in particular – in populations with [unexplained physical symptoms](#).

PTSD is associated with [changes in brain function](#), hormonal imbalance, and [immune system](#) dysregulation. Hormonal dysregulation, particularly involving stress hormones like cortisol and adrenaline, leads to heightened arousal and chronic stress, compromising immune function and increasing susceptibility to infections and inflammation.

A study published in Lancet Psychiatry indicates that PTSD [elevates the risk of cardiac events up to 27%](#). Chronic stress contributes to high blood pressure, while inflammation exacerbates chronic pain conditions like fibromyalgia. Unhealthy coping behaviors associated with PTSD, such as smoking and excessive alcohol consumption, further increase health risks.

## **Specialized Care and Treatment Options for PTSD**

PTSD is a highly individualized disorder and requires personalized care approaches. No two cases are the same, and no two treatments will be the same. [Trauma-informed care](#), now considered the gold standard across healthcare settings, involves thoroughly examining a patient's entire situation, including any past traumatic experiences.

## **Importance of Specialized Care and Treatment**

Specialized care combines proven methods with personalized approaches. Patients often see significant improvements with specialized care. From fewer PTSD symptoms to better ways of coping, it's all about feeling understood and supported by your healthcare team.

The treatment landscape for PTSD offers a diverse array of options, underlining the importance of exploring various therapeutic modalities. The goal of these treatments is to alleviate symptoms, foster healing, and enhance the overall quality of life for individuals dealing with PTSD.

## **Therapies for PTSD**

### **Cognitive Behavioral Therapy (CBT)**

[CBT](#) combines various techniques to address both the cognitive and behavioral aspects of PTSD, empowering individuals to challenge negative beliefs, develop

effective coping strategies, and regain control over their lives in the aftermath of trauma.

CBT is strongly recommended for the [treatment of PTSD](#) and uses the following methods:

- Psychoeducation
- Assessment and goal-setting
- Cognitive restructuring
- Exposure therapy
- Skills training
- Relapse prevention

Some examples of techniques used in CBT for PTSD include:

- Creating a trauma narrative
- Gradual exposure
- Grounding techniques
- Cognitive coping strategies

### **Cognitive Processing Therapy (CPT)**

Delivered over 12 sessions and rooted in CBT-like education and relapse prevention, [CPT](#) provides a structured approach to addressing PTSD symptoms by targeting maladaptive cognitive processes, facilitating emotional processing, and enhancing coping skills.

Clients are encouraged to confront and process distressing thoughts and feelings associated with the trauma within the safety of the therapeutic environment. CPT focuses on processing through structured writing exercises known as cognitive worksheets or cognitive processing worksheets. These worksheets guide individuals through exploring and challenging their thoughts and beliefs about the traumatic event. The goal is to create a coherent narrative of the trauma and its aftermath while addressing cognitive distortions.

### **Cognitive Therapy (CT)**

[CT](#) is a therapeutic approach focused on identifying and modifying thought patterns and beliefs that contribute to emotional distress. CT is effective in the treatment of PTSD, as it targets the cognitive aspects of the disorder, helping individuals

challenge and change their perceptions and interpretations of traumatic events.

The framework for using CT for PTSD includes the following steps:

- **Identifying cognitive distortions**

Common distortions include [catastrophizing](#), [overgeneralization](#), and [personalization](#).

- **Challenging negative automatic thoughts**

Clients are encouraged to question the accuracy and validity of these thoughts and consider alternative, more balanced perspectives.

- **Examining core beliefs**

Core beliefs may include notions of safety, trust, and self-worth that have been profoundly affected by the traumatic experience.

- **Reconstructing trauma narratives**

This process allows individuals to integrate new information and perspectives, reducing the emotional intensity and power of the trauma memory.

## **Prolonged Exposure Therapy (PE)**

PE is a [highly effective treatment](#) for PTSD that helps individuals confront and process traumatic memories through exposure-based techniques.

[PE](#) therapy for PTSD operates on two main principles:

1. Individuals with PTSD often avoid reminders of their trauma, exacerbating distress and impairing functioning. PE disrupts this avoidance pattern by facilitating the emotional processing of traumatic memories.
2. Through repeated exposure to trauma-related stimuli in a safe environment, PE aims to habituate individuals to these triggers and extinguish conditioned fear responses associated with the trauma.

Concerns about PE therapy for PTSD are common. Despite fears of re-traumatization, [evidence shows](#) PE's safety and efficacy. Therapists collaborate with clients to ensure exposure exercises progress at a manageable pace. The [National Center for PTSD](#) states, "PE is typically delivered in 8 to 15, 90-minute sessions, weekly."

## **Conditionally Recommended Therapies**



Conditionally recommended therapies for PTSD offer alternative approaches for individuals who may not fully respond to the first-line treatments mentioned above.

### **Brief Eclectic Psychotherapy (BEP)**

[BEP](#) is a [time-limited therapy](#) that integrates elements from various therapeutic modalities, including cognitive-behavioral, psychodynamic, and humanistic approaches. It spans 16 sessions, each lasting 45 to 60 minutes and with specific objectives.

BEP involves a collaborative and individualized treatment approach, where therapists work with clients to identify and address specific cognitive, emotional, and interpersonal issues related to their trauma. BEP works best with people experiencing [PTSD from one traumatic event](#). Techniques may include cognitive restructuring, emotion regulation skills, and exploring underlying relational patterns.

While research on BEP is limited, one older [study](#) suggests it may reduce PTSD symptoms by targeting both cognitive and emotional aspects of the disorder.

### **Eye Movement Desensitization and Reprocessing (EMDR)**

[Numerous studies](#) support EMDR's effectiveness in reducing PTSD symptoms and facilitating trauma processing. It's shown to be as effective as traditional therapies like [CPT](#) and [PE](#), with rapid and lasting improvements observed in some individuals after just a few sessions.

[EMDR](#) involves bilateral stimulation, such as eye movements, to facilitate the processing of traumatic memories. An EMDR session typically proceeds as follows:

- **History-Taking:** Gathering information about trauma history and treatment goals.
- **Preparation:** Providing psychoeducation and teaching coping skills.
- **Desensitization:** Focusing on a targeted traumatic memory while engaging in bilateral stimulation.
- **Reprocessing:** Facilitating the reprocessing of the traumatic memory and integrating new beliefs.
- **Closure:** Ensuring stability and grounding at the end of each session.
- **Reevaluation:** Reviewing progress and identifying new targets for processing.

### **Narrative Exposure Therapy (NET)**

[NET](#) is a [short-term therapy](#) designed to address PTSD symptoms by systematically recounting and processing traumatic memories within the context of a client's life story.

NET involves a structured series of sessions to construct a detailed timeline of the client's life, including traumatic events and associated emotions and cognitions. Therapists guide clients through narrating their experiences, gradually exposing them to distressing memories while providing support and validation.

NET has demonstrated [promising results in reducing PTSD](#) symptoms across various populations, including refugees, survivors of mass violence, and individuals with complex trauma histories. [Research](#) suggests that NET may be particularly effective in populations where traditional exposure-based therapies are challenging to implement.

## Medication-Based Treatment

The [2023 Clinical Practice Guidelines for PTSD](#) urge psychotherapy to be the first choice in treatment for PTSD. The guidelines discuss that medication can be an effective treatment alongside therapy or when therapies aren't available, and a patient has requested help from medication. In such cases, medication can play a significant role in the treatment of PTSD, particularly in managing comorbid symptoms such as anxiety, depression, and insomnia that often accompany the disorder.

While medications do not address the underlying trauma, they can help alleviate distressing symptoms and improve overall functioning. It's essential to have open discussions with healthcare providers to assess the benefits and risks of medication.

The following table shows classes of medications commonly used, their generic and trade name, and mechanism of action. It's important to note that of the below medications, currently, only SSRIs are [FDA-approved](#) for PTSD. Other medications noted for PTSD are for off-label use.

Drug Class	Generic Name	Trade Name	How it works
------------	--------------	------------	--------------

Selective Serotonin Reuptake Inhibitors (SSRIs)	<ul style="list-style-type: none"> <li>• Sertraline</li> <li>• Paroxetine</li> <li>• Fluoxetine</li> </ul>	<ul style="list-style-type: none"> <li>• Zoloft</li> <li>• Paxil</li> <li>• Prozac</li> </ul>	Increase the levels of <a href="#">serotonin</a> in the brain, which can help regulate mood and reduce anxiety and depression symptoms.
Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)	<ul style="list-style-type: none"> <li>• Venlafaxine</li> <li>• Duloxetine</li> </ul>	<ul style="list-style-type: none"> <li>• Effexor</li> <li>• Cymbalta</li> </ul>	Increase serotonin levels in the brain but additionally target <a href="#">norepinephrine</a> .
Benzodiazepines	<ul style="list-style-type: none"> <li>• Clonazepam</li> <li>• Lorazepam</li> </ul>	<ul style="list-style-type: none"> <li>• Klonopin</li> <li>• Ativan</li> </ul>	Enhances the effects of <a href="#">GABA</a> , leading to sedation, muscle relaxation, and anxiety reduction. Can be habit-forming.
Prazosin	<ul style="list-style-type: none"> <li>• Prazosin</li> </ul>	<ul style="list-style-type: none"> <li>• Minipress</li> </ul>	It can help reduce the frequency and intensity of nightmares and improve sleep quality.

## Integrative and Holistic Approaches

Integrative and holistic approaches to treating PTSD incorporate complementary therapies alongside traditional treatments to address the interconnectedness of mind, body, and spirit for comprehensive recovery. By integrating complementary therapies, individuals can access additional tools and resources to support their well-being.

- **Mindfulness-based interventions:** Practices like mindfulness meditation and yoga [reduce symptoms](#) by promoting relaxation, alleviating symptoms, and emotional regulation.
- **[Complementary therapies:](#)** Practices such as acupuncture and massage therapy can reduce stress and promote relaxation.

## Supportive Services for PTSD

Supportive services are essential in the recovery journey of individuals with PTSD, providing emotional support, practical assistance, and fostering resilience. They include support groups, online resources, family assistance programs, and

community organizations, offering peer connection, educational resources, and access to mental health resources. These services empower individuals to navigate the challenges of PTSD and build a supportive network for healing and recovery.

## **Support Groups and Communities**

The understanding, empathy, and support offered by peers can make a significant difference in navigating the challenges of PTSD and building a solid support network for long-term well-being.

Here's why participation in these groups is highly encouraged:

- **Peer support:** Interacting with peers with similar experiences fosters a sense of validation, understanding, and belonging.
- **Shared experiences:** In support groups, individuals can share their stories, challenges, and successes in a nonjudgmental environment.
- **Mutual understanding:** Support group members can empathize with each other's struggles in ways that friends and family may not fully understand.

If you're currently receiving therapy for PTSD, ask your therapist or mental health provider for recommendations on local support groups or online communities tailored to individuals with PTSD.

## **Online Services and Resources**

Online services and resources offer convenient and accessible support for individuals with PTSD. Here's why they're beneficial:

- **24/7 availability:** Online resources are accessible anytime, anywhere, allowing individuals to seek support and information conveniently.
- **Anonymity:** Many online platforms offer anonymity, making it easier for individuals to seek help without fear of judgment or stigma.
- **Wide range of resources:** From informational websites to self-help tools and virtual support groups, online services provide a diverse range of resources to meet the varied needs of individuals with PTSD.

## **Reputable Online Resources**

[The National Center for PTSD](#)

- Provides information on evidence-based treatments, self-help resources, and links to find specialized care providers.

#### [The Anxiety and Depression Association of America](#)

- Offers resources and a therapist directory for finding mental health professionals specializing in PTSD treatment.

#### [National Institute of Mental Health: PTSD](#)

- Detailed information on the ins and outs of PTSD, with treatment options, clinical trial information, and added resources.

#### [Veterans Crisis Line](#)

- 24/7, confidential crisis support for you or a loved one.

#### [Substance Abuse and Mental Health Services Administration](#)

- Raising awareness of substance abuse with multiple crisis lines, funding opportunities, and resources to find treatment.

### **Family and Social Support**

Support from family and friends is indispensable for individuals dealing with PTSD. They offer emotional support through compassionate listening, validation, and empathy. Practical assistance with daily tasks like household chores or childcare helps alleviate stress, allowing the individual to focus on recovery.

Guidance for loved ones includes educating themselves about PTSD, being patient and nonjudgmental, offering practical help while respecting boundaries, and encouraging professional treatment. Through a combination of emotional support, practical assistance, and understanding, family and friends can significantly contribute to the recovery journey of individuals with PTSD.

### **Challenges in PTSD Care**

Providing adequate care for individuals with PTSD is often hindered by various challenges that can impede access to treatment and hinder recovery.

### **Limitations and Obstacles in PTSD Treatment**

Current PTSD treatment approaches face several limitations and obstacles that can impact treatment outcomes and accessibility. Several factors contribute to the resistance or ineffectiveness of PTSD treatment. These include, but are not limited to:

- **Variability in treatment response:** Not all individuals respond to traditional PTSD treatments, leading to variability in treatment outcomes.
- **High dropout rates:** Some individuals may discontinue treatment prematurely due to factors such as lack of motivation, dissatisfaction with treatment progress, or difficulty tolerating treatment-related distress. The presence of comorbid conditions, such as depression, substance use disorders, or personality disorders, can complicate treatment efforts and contribute to resistance.
- **Access barriers:** Limited access to mental health services, long wait times for appointments, and financial constraints can hinder individuals from receiving timely and appropriate care.

These factors collectively underscore the complexities involved in addressing PTSD and highlight the importance of comprehensive and tailored approaches to treatment.

### **Overcoming Barriers to Accessing PTSD Care**

Accessing care for PTSD can be daunting due to a variety of barriers, including geographic limitations, financial constraints, stigma, and cultural factors. Many individuals find themselves in areas with limited mental health resources, particularly in rural or remote regions, making it difficult to access appropriate care. Financial barriers, such as lack of insurance coverage or high out-of-pocket costs, can pose significant obstacles to seeking treatment. Cultural factors, including language barriers and a lack of culturally responsive services, further compound the challenge of accessing PTSD care.

To overcome these barriers, individuals can explore [telehealth](#) options, which provide remote access to mental health services. Checking insurance coverage and seeking out mental health providers who offer sliding-scale fees based on income can also make therapy more affordable for those with financial limitations.

[Community-based programs](#), [nonprofit organizations](#), and [government-funded initiatives](#) often offer low-cost or free mental health services.

Advocacy for policy changes is also crucial in improving access to PTSD care. Supporting policy initiatives that prioritize mental health, increase funding for mental health services, and promote culturally responsive care for underserved populations can help address systemic barriers to care.

## **Addressing Stigma Associated With PTSD**

The [stigma surrounding PTSD](#) can have profound effects on individuals' willingness to seek help, engage in treatment, and integrate into social environments. Misconceptions and stereotypes about PTSD often lead to fear, shame, and discrimination. This [stigma](#) can result in delayed or avoided help-seeking, as individuals may fear judgment or believe that seeking help is a sign of weakness.

Challenging these misconceptions and stereotypes is essential in combating stigma and fostering a supportive environment for individuals with PTSD. By sharing accurate information about PTSD, we can dispel myths and educate others about the realities of the disorder.

[Personal stories](#) of resilience and examples of successful treatment outcomes can also challenge stigma by highlighting the strength and courage of individuals living with PTSD. These narratives demonstrate that recovery is possible, and that seeking help is a brave and empowering choice.

## **Future Perspective on PTSD Care**

As we look to the future, the outlook for PTSD care is promising, with potential advancements on the horizon that could significantly improve treatment efficacy, accessibility, and societal understanding of the disorder.

## **Advances in PTSD Treatment**

Recent advances in evidence-based treatments for PTSD have broadened treatment options, including novel psychotherapeutic approaches like [CPT](#) and [PE](#). Pharmacological interventions such as SSRIs and SNRIs are commonly prescribed, with emerging research exploring the potential of adjunctive treatments like [ketamine and MDMA](#). Integrative modalities like mindfulness-based interventions and yoga are gaining recognition for their holistic approach.

Challenges persist in translating research into real-world applications, including disparities in access to treatment and the need for tailored multidisciplinary

approaches. Continued collaboration between researchers, clinicians, and policymakers is essential to address these challenges and improve outcomes for individuals with PTSD.

## **The Promise of Technology in PTSD Care**

Advancements in technological care for PTSD have the potential to expand access to care and reach underserved populations. Digital literacy among patients and providers is crucial to utilize technology-based interventions effectively. Healthcare professionals and policymakers must address disparities in access to digital resources, particularly among underserved populations facing barriers such as limited Internet connectivity or lack of technological proficiency.

Some exciting prospects in technology for PTSD include:

- [Virtual reality \(VR\)](#) simulations enable exposure therapy in a controlled and immersive environment, allowing individuals to confront and process traumatic memories in a safe setting.
- [Mobile apps](#) offer self-help tools, symptom tracking, and psychoeducation, empowering individuals to manage their symptoms and access support anytime, anywhere.
- [Wearable devices](#) provide valuable insights for individuals and clinicians in understanding symptom patterns and identifying triggers.
- [Telemedicine](#) platforms help facilitate remote therapy sessions.

## **Encouraging a More Compassionate Society for PTSD**

Education plays a crucial role in raising awareness about PTSD, its symptoms, and its impact on individuals' lives. By providing accurate information and dispelling myths and misconceptions surrounding the disorder, education helps reduce stigma and foster empathy among the general public.

Trauma-informed care training for healthcare professionals, first responders, and educators is essential to deliver services in a sensitive and supportive manner. Campaigns like [PTSD Awareness Month](#) in June and events like [PTSD Awareness Day](#) on June 27 help raise public awareness. Education and awareness are vital to promote understanding and contribute to creating a more compassionate and supportive society for individuals living with PTSD.



We hope this article has given you the education you need to seek treatment for your PTSD. There are people out there who want to help, and there are many resources available.

[Amanda Farquharson, BSN, RN](#)

Updated May 7, 2024

Reviewed by [NursingEducation Staff](#)

- [FAQ](#)

[Missouri Lawyers Corner Veterans](#)

Print

Table of Contents

NEWS

News & publications

The news about recent activities for needed peoples.

[More News](#)

28 Jul 2025



See what cuts to LSC would mean to your community

Actions speak louder than words. Last week, members of Congress sent a powerful...

[Continue Reading about See what cuts to LSC would mean to your community](#)

18 Jun 2025



Key Groups Send Support Letters to Congress for LSC

Amid threats to LSC's funding, key groups are stepping up to urge Congress's...

[Continue Reading about Key Groups Send Support Letters to Congress for LSC](#)

---

*PDF downloaded from <https://www.lsmo.org/node/1062/ptsd-specialized-care-and-supportive-services>*